

ALCOHOL TALLY SHEET FOR SECTION E

DSM-III-R

.....E9 3+ times wanted to stop/cut down
.....E9.C Found was unable to stop/cut down
GROUP 1: _____

.....E12.A (E12.B) Neglected responsibilities when bingeing 3+ times
.....E23 Drinking/being hung over often interfered with responsibilities
.....E27.C Arrested for drunk driving/had accidents/damaged car 3+ times
.....E29.B Alcohol caused accidental injuries 3+ times
.....E30.B Drunk in situations where could have injured self 3+ times
.....E36.C Mixed alcohol & drugs 3+ times when knew this was dangerous
GROUP 2: _____

.....E13.C Drank more than intended/more days in row or when
promised self wouldn't 3+ times
.....E14.B Drunk when didn't want to 3+ times
GROUP 3: _____

.....E15 While drinking/recovering from effects/little time for anything else.
GROUP 4: _____

.....E17.2 Drinking caused problems with family/friends 3+ times
.....E17.3 Lost friends because of drinking 3+ time
.....E17.4 Drinking caused problems with work/school 3+ times
.....E24.B Drank knowing it caused problems with love relationships
.....E28.B Arrested/detained by police for drunk behavior 3+ times
.....E35.B Continued to drink knowing alcohol caused health problems
.....E36 Continued to drink despite serious illness
.....E37.A Continued to drink knowing alcohol caused emotional problems
GROUP 5: _____

.....E20.A Needed increase in drinking of 50%
.....E20.C Could drink 50% more before drunk
GROUP 6: _____

.....E22 Given up/greatly reduced important activities to drink
GROUP 7: _____

.....E32 Experienced withdrawal symptoms (III-R requires shakes + 1)
.....E33 Had fits/seizures from drinking
.....E34 Had the DT's from drinking
GROUP 8: _____

.....E32.H 3+ occasions drank to relieve or avoid withdrawal symptoms
.....E33.D Drank to relieve/avoid fits/seizures
.....E34.D Drank to relieve/avoid the DT's 3+ times
GROUP 9: _____

ALCOHOL TALLY SHEET FOR SECTION E

ICD-10

_____.....E11 Strong desire for drink

GROUP 1: _____

_____.....E9.C Found was unable to stop/cut down
_____.....E11 Drank when promised self not to/drank more than intended
_____.....E13.A Drank more days in row than intended
_____.....E14 Became drunk when didn't want to

GROUP 2: _____

_____.....E16 Drinking became regular/wouldn't change no matter what

GROUP 3: _____

_____.....E20 Needed more alcohol to get an effect
_____.....E20.B Found could drink a lot more before drunk

GROUP 4: _____

_____.....E22 Given up/greatly reduced important activities to drink

GROUP 5: _____

_____.....E29.B Alcohol caused accidental injuries 3+ times
_____.....E35.B Continued to drink knowing alcohol caused health problems
_____.....E36 Continued to drink despite serious illness
_____.....E37.A Continued to drink knowing alcohol caused emotional problems

GROUP 6: _____

_____.....E32 Experienced withdrawal symptoms (shakes + 1)
_____.....E33 Had fits/seizures from drinking
_____.....E34 Had the DT's from drinking

GROUP 7: _____

_____.....E32.H 3+ occasions drank to relieve or avoid withdrawal symptoms
_____.....E33.D Drank to relieve/avoid fits/seizures
_____.....E34.D Drank to relieve/avoid the DT's 3+ times

GROUP 8: _____

C O G A/SSAGA

MARIJUANA TALLY SHEET FOR SECTION F

DSM-IIIIR

_____.....F5 Great deal of time spent using marijuana, getting it or getting over its effects
GROUP 1: _____

_____.....F7 Often wanted to stop/cut down on marijuana
_____.....F7.A Found was unable to stop/cut down on marijuana
GROUP 2: _____

_____.....F8 Often used marijuana more frequently or in larger amounts than intended
GROUP 3: _____

_____.....F9 Needed larger amounts of marijuana to get same effect/couldn't get high on usual amount
GROUP 4: _____

_____.....F10 Experienced withdrawal from cutting down/stopping marijuana
GROUP 5: _____

_____.....F10.A 3+ times used marijuana to relieve/avoid withdrawal symptoms
GROUP 6: _____

_____.....F11.A 3+ times high from marijuana when could have injured self
_____.....F11.C Marijuana 3+ times caused accidental injuries
_____.....F14 Being high/having withdrawal often interfered with responsibilities
GROUP 7: _____

_____.....F6.A Continued to use marijuana knowing it caused emotional or psychological problems, or decreased contact with family/friends
_____.....F12.A Continued to use marijuana despite problems with family/friends
GROUP 8: _____

_____.....F13 3+ times gave up/greatly reduced important activities to use marijuana
GROUP 8: _____

C O G A/SSAGA

MARIJUANA TALLY SHEET FOR SECTION F

ICD-10

_____.....F6.A Continued to use marijuana knowing it caused emotional or psychological problems, or decreased contact with family/friends
_____.....F11.C Marijuana 3+ times caused accidental injuries
GROUP 1: _____

_____.....F7.A Found was unable to stop/cut down on marijuana
_____.....F8 Often used marijuana more frequently or in larger amounts than intended
GROUP 2: _____

_____.....F9 Needed larger amounts of marijuana to get same effect/couldn't get high on usual amount
GROUP 3: _____

_____.....F10 Experienced withdrawal from cutting down/stopping marijuana
GROUP 4: _____

_____.....F10.A 3+ times used marijuana to relieve/avoid withdrawal symptoms
GROUP 5: _____

_____.....F13 3+ times given up/greatly reduced important activities to use marijuana
GROUP 6: _____

_____.....F15 Strong desire for marijuana/couldn't think of anything else
GROUP 7: _____

_____.....F16 Marijuana use became so regular wouldn't change no matter what
GROUP 8: _____

C O G A/SSAGA

DSM-IIIIR DRUG TALLY SHEET FOR SECTION G

		COCAINE	STIM.	SED.	OPIATE	OTHER
G6	A month or more spent using, getting, or getting over effects of (DRUG)	_____	_____	_____	_____	_____
G7	Often wanted to cut down on (DRUG) or tried to stop/cut down on (DRUG) but couldn't	_____	_____	_____	_____	_____
G8	Needed larger amounts of (DRUG) to get effect or couldn't get high on same amount	_____	_____	_____	_____	_____
G9	Gave up/reduced important activities to use (DRUG)	_____	_____	_____	_____	_____
G10	Used (DRUG) more days or in larger amounts than meant to	_____	_____	_____	_____	_____
G11.B	Experienced withdrawal from (DRUG)	_____	_____	_____	_____	_____
G11.D	Used (DRUG) 3+ times to relieve or avoid withdrawal symptoms	_____	_____	_____	_____	_____
G12.B1	(DRUG) 3+ times caused an overdose ..	_____	_____	_____	_____	_____
G12.C1	Continued to use (DRUG) knowing it caused hepatitis	_____	_____	_____	_____	_____
G12.D1	Continued to use (DRUG) knowing it caused other health problems	_____	_____	_____	_____	_____
G13.E	Continued to use (DRUG) knowing it caused problems/objections from family/others, problems at work/school, physical fights, trouble with police	_____	_____	_____	_____	_____
G15.A	Continued to use (DRUG) knowing it caused emotional/psychological problems	_____	_____	_____	_____	_____
G12.A1	(DRUG) 3+ times caused accidental injuries	_____	_____	_____	_____	_____
G14	Being high on (DRUG)/suffering its after-effects often interfered with working/school/other responsibilities	_____	_____	_____	_____	_____
G16	3+ times high on (DRUG) when could have gotten hurt	_____	_____	_____	_____	_____

ICD-10 DRUG TALLY SHEET FOR SECTION G

		COCAINE	STIM.	SED.	OPIATE	OTHER
G4	Use of (DRUG) became so regular wouldn't change no matter what	_____	_____	_____	_____	_____
G5	Strong desire for (DRUG) that hard to think of anything else	_____	_____	_____	_____	_____
G7	Often wanted to cut down on (DRUG) or tried to stop/cut down on (DRUG) but couldn't	_____	_____	_____	_____	_____
G10	Used (DRUG) more days or in larger amounts than meant to	_____	_____	_____	_____	_____
G8	Needed larger amounts of (DRUG) to get effect or couldn't get high on same amount	_____	_____	_____	_____	_____
G9	Gave up/reduced important activities to use (DRUG)	_____	_____	_____	_____	_____
G11.B	Experienced withdrawal from (DRUG)	_____	_____	_____	_____	_____
G11.D	Used (DRUG) 3+ times to relieve or avoid withdrawal symptoms	_____	_____	_____	_____	_____
G12.A1	(DRUG) 3+ times caused accidental injuries	_____	_____	_____	_____	_____
G12.B1	(DRUG) 3+ times caused an overdose ..	_____	_____	_____	_____	_____
G12.C1	Continued to use (DRUG) knowing it caused hepatitis	_____	_____	_____	_____	_____
G12.D1	Continued to use (DRUG) knowing it caused other health problems	_____	_____	_____	_____	_____
G15.A	Continued to use (DRUG) knowing it caused emotional/psychological problems	_____	_____	_____	_____	_____

TALLY SHEET FOR SECTION M
(QUESTION M38)

CHECK ONLY IF BEHAVIOR OCCURED AFTER 15TH BIRTHDAY
(COUNT ITEMS CODED 5 OR 6)

- M5 (M5B,D,F) Stealing/forging signature_____
- M6 (M6C) Damaging property_____
- M7 Starting fights_____
- M7B Getting into physical fights_____
- M8 Using a weapon in a fight_____
- M9 Purposely injuring someone_____
- M14 Being a bully_____
- M15 Being mean to animals_____
- M16 Setting fires_____
- M17 Outsmarting people in authority_____
- M18 Breaking and entering_____
- M19 Mugging/armed robbery_____
- M21.1 Writing bad checks_____
- M21.2 Selling drugs receiving/buying stolen goods..._____
- M21.5 Finding customers for prostitutes_____
- M21.A Other arrestable behavior_____
- M24.A Never being faithful_____
- M26 Forced others into sexual activity_____
- M28 Hitting or assaulting family/non-family_____
- M33 Defaulting on debts_____
- M34 Not caring for children when supposed to_____